



**MAY 2012**

## **THE WINTER-RUN HARVEST MODEL (WRHM)**

Michael R. O'Farrell, Shanae D. Allen, and Michael S. Mohr

NOAA-TM-NMFS-SWFSC-489

U.S. DEPARTMENT OF COMMERCE  
National Oceanic and Atmospheric Administration  
National Marine Fisheries Service  
Southwest Fisheries Science Center

The National Oceanic and Atmospheric Administration (NOAA), organized in 1970, has evolved into an agency that establishes national policies and manages and conserves our oceanic, coastal, and atmospheric resources. An organizational element within NOAA, the Office of Fisheries is responsible for fisheries policy and the direction of the National Marine Fisheries Service (NMFS).

In addition to its formal publications, the NMFS uses the NOAA Technical Memorandum series to issue informal scientific and technical publications when complete formal review and editorial processing are not appropriate or feasible. Documents within this series, however, reflect sound professional work and may be referenced in the formal scientific and technical literature.



## NOAA Technical Memorandum NMFS

This TM series is used for documentation and timely communication of preliminary results, interim reports, or special purpose information. The TMs have not received complete formal review, editorial control, or detailed editing.

**MAY 2012**

# THE WINTER-RUN HARVEST MODEL (WRHM)

Michael R. O'Farrell<sup>1</sup>, Shanae D. Allen<sup>1,2</sup>, and Michael S. Mohr<sup>1</sup>

<sup>1</sup>National Oceanic & Atmospheric Administration  
National Marine Fisheries Service, NOAA  
Southwest Fisheries Science Center  
Fisheries Ecology Division

<sup>2</sup>Institute of Marine Science; University of California, Santa Cruz

NOAA-TM-NMFS-SWFSC-489

### U.S. DEPARTMENT OF COMMERCE

Rebecca M. Blank, Secretary

**National Oceanic and Atmospheric Administration**

Jane Lubchenco, Undersecretary for Oceans and Atmosphere

**National Marine Fisheries Service**

Eric C. Schwaab, Assistant Administrator for Fisheries



# 1 Abstract

The Winter Run Harvest Model (WRHM) forecasts the annual age-3 ocean impact rate on Sacramento River winter Chinook resulting from salmon fisheries south of Point Arena, California. This impact rate includes both landed and non-landed mortality attributable to fisheries. The model is a tool developed for use in the Pacific Fishery Management Council (PFMC) arena for managing fisheries to comply with the National Marine Fisheries Service Endangered Species Act consultation standard for Sacramento River winter Chinook beginning in 2012. Analogous to other models used for assessment and management of salmon through the PFMC process, the WRHM is temporally and spatially stratified. Impact rates are forecast for each month, area, and sector (commercial, recreational) to capture variation in exploitation patterns and fishery management measures that occur at that scale. A forecast of the total age-3 impact rate is then made by aggregating impacts over all strata where fishing occurred. The WRHM is capable of accounting for the customary fishery management measures used by the PFMC (e.g., time/area/sector closures, quotas, and minimum size limits) and therefore readily integrates into the annual PFMC salmon fishery planning process.


## 2 Introduction

Sacramento River winter Chinook (SRWC) is an endangered Chinook salmon (*Oncorhynchus tshawytscha*) stock harvested incidentally in ocean fisheries. First listed under the Endangered Species Act (ESA) as threatened in 1989, SRWC status was downgraded to endangered in 1994. Most recently, in the 2010 Biological Opinion for ocean fisheries (NMFS 2010), the National Marine Fisheries Service (NMFS) found that ocean fisheries are likely to jeopardize the continued existence of SRWC owing to a lack of measures and tools to constrain or reduce fishery impacts when SRWC population status is poor. NMFS offered a reasonable and prudent alternative (RPA) to comply with the ESA, which included (1) establishing thresholds related to the status of SRWC, (2) establishing fishery management objectives, and (3) development of analytical tools and assessment models that can implement the fishery management objectives in the salmon fishery management process. This report documents one portion of component (3): the Winter Run Harvest Model (WRHM).

To implement the RPA specified in the 2010 Biological Opinion, a two-part SRWC consultation standard was developed. The first component of the consultation standard is to continue requiring the time/area/fishery closure and size limit provisions identified in the 2004 Biological Opinion each year, regardless of SRWC status (NMFS 2004):

The recreational salmon fishery between Point Arena and Pigeon Point shall open no earlier than the first Saturday in April and close no later than the second Sunday in November. The recreational salmon fishery between Pigeon Point and the U.S.–Mexico Border shall open no earlier than the first Saturday in April and close no later than the first Sunday in October. The minimum size limit shall be at least 20 inches total length.

The commercial salmon fishery between Point Arena and the U.S.–Mexico border shall open no earlier than May 1 and close no later than September 30, with the exception of an October fishery conducted Monday through Friday between Point Reyes and Point San Pedro, which shall end no later than October 15. The minimum size



**Figure 1.** Sacramento River winter Chinook fishery control rule.

limit shall be at least 26 inches total length.


The WRHM was developed to implement the second component of the SRWC consultation standard: a control rule that limits the maximum age-3 impact rate (allowable as a preseason forecast) based on the geometric mean of the most recent three years of spawner escapement (Figure 1). When the three-year geometric mean of spawner escapement is in excess of 5000, a maximum forecast age-3 impact rate is not specified and the consultation standard reduces to only the first component. When the three-year geometric mean is between 4000 and 5000, the maximum forecast age-3 impact rate is 0.20. Between 3-year geometric mean values of 4000 and 500, the maximum forecast age-3 impact rate decreases linearly from 0.20 to 0.10. Finally, at 3-year geometric mean spawner levels less than 500, the maximum forecast age-3 impact rate is zero. The age-3 impact rate specified by the control rule, and forecast by the WRHM, includes both landed and non-landed mortality attributable to fisheries, and the region covered includes the San Francisco (SF) management area (Point Arena to Pigeon Point) and the Monterey (MO) management area (Pigeon Point to the US/Mexico border) in California.

For SRWC, the age-3 ocean fishery impact rate is an appropriate metric for use in controlling overall fisheries exploitation. The age-3 impact rate closely approximates the cohort's spawner reduction rate, which is the fraction of a cohort's potential spawners that are eliminated by the fishery (see Figure 5 in O'Farrell et al. 2012). The concordance between the age-3 impact rate and the spawner reduction rate is due to the very high (> 85 percent) age-3 maturation rates SRWC exhibit (O'Farrell et al. 2012). In addition, the age-3 impact rate can be forecast in the absence of a SRWC preseason abundance forecast. A preseason abundance forecast cannot be made for SRWC in time for the PFMC preseason management process due to the timing of SRWC spawning and the timing of annual ocean salmon fisheries. The age-2 (jack) river return data that would be necessary to forecast age-3 abundance prior to spring/summer ocean salmon fisheries are not available until the fall or winter following those fisheries and therefore are not useful for making a timely age-3 ocean abundance forecast.

Forecasts of the age-3 impact rate will be confined to fisheries occurring in management areas south of Point Arena because the overwhelming majority of SRWC impacts occur in this region. Figure 2 demonstrates that in most years for which the age-3 impact rate has been estimated, zero impacts resulted from fisheries north of Point Arena, and when they did occur, they represented a very small portion of the overall age-3 impact rate. Between years 2000 and 2007, the age-3 impact rate attributed to fisheries north of Point Arena averaged 0.0058.

The WRHM consists of projecting an age-3 cohort abundance through ocean fisheries on a monthly basis between the beginning of March 1 (year  $y$ ) and the end of the last day of February ( $y + 1$ ). The starting abundance is arbitrary and does not affect the forecast of the annual age-3 impact rate; hereafter we assume that the March 1 ( $y$ ) ocean abundance is equal to 1.0. March 1 was chosen as the "birth date" for SRWC, based on the reported peak migration period into the Sacramento River basin from Fisher (1994). Monthly age-3 impacts, forecast by area (SF, MO), and sector (commercial, recreational) under the proposed fishery management measures, are deducted from the monthly abundance. The total age-3 impact rate is then computed by totaling the month/area/sector impacts and dividing by the assumed March 1 ( $y$ ) ocean abundance. The WRHM is able to accommodate days-open (fisheries specified as the number of days open to fish-





**Figure 2.** Age-3 ocean fishery impact rate, partitioned by contributions from fisheries north and south of Point Arena, CA.

ing and not as a harvest limit) and quota fishery management measures, with one exception that is explained in more detail in section 4.2. The WRHM is also able to account for variation in minimum size limits. Hence, management measures such as month/area/sector closures and minimum size limits commonly used by the PFMC to constrain the salmon fishery can be directly accounted for in the WRHM-derived forecast of the age-3 impact rate.

Documentation of the WRHM follows in sections 3 and 4. Section 3 defines the main model structure, methods used to project the age-3 cohort through ocean fisheries, and the expression used to forecast the age-3 impact rate. Section 4 describes the submodels and input variables used to parameterize the WRHM. The report ends with a discussion of key components of the model and a comparison to existing PFMC harvest models for Chinook salmon.

### 3 Main model

The age-3 SRWC cohort abundance is projected through ocean fisheries sequentially from  $t =$  March ( $y$ ) through  $t =$  February ( $y + 1$ ). The method of forward projection of the age-3 cohort

**Table 1.** Notation used in this memorandum.

Symbol	Definition
$\bar{\phantom{x}}$	Overbar denoting the arithmetic mean
$\beta$	Contact rate per unit effort
$C$	Ocean fishery contacts
$c$	Contact rate
com	Term denoting the commercial fishery
$D$	Number of deaths due to drop off mortality
$d$	Drop off mortality rate
$f$	Fishing effort
$H$	Harvest
$I$	Impacts
$i$	Impact rate
$l$	Total length (inches)
$l^*$	Minimum size limit for ocean fisheries; total length (inches)
MO	Monterey major port area
$\mu$	Mean length
$N$	Cohort abundance
$p$	Proportion of cohort abundance expected to be $\geq$ the minimum legal size
rec	Term denoting the recreational fishery
$\rho$	Proportion of the recreational Chinook harvest caught by mooching
$S$	Number of deaths due to release mortality
$s$	Release mortality rate
SF	San Francisco major port area
$\sigma$	Standard deviation of length
$t$	Subscript denoting month
$v$	Monthly natural mortality rate
$x$	Subscript denoting sector, $x \in \{\text{com,rec}\}$
$y$	Subscript denoting year
$z$	Subscript denoting area, $z \in \{\text{SF,MO}\}$

is consistent with the backward reconstruction of cohorts described for SRWC in O'Farrell et al. (2012). Notation for all variables and parameters used in this report are found in Table 1.

For each month  $t$ , the following metrics are computed by area  $z$  and sector  $x$ :

$$C_{tzx} = c_{tzx} \times N_t \quad (1)$$

$$H_{tzx} = C_{tzx} \times p_{tzx} \quad (2)$$

$$S_{tzx} = (C_{tzx} - H_{tzx}) \times s_{tzx} \quad (3)$$

$$D_{tzx} = C_{tzx} \times d \quad (4)$$

$$I_{tzx} = H_{tzx} + S_{tzx} + D_{tzx}, \quad (5)$$

with cohort abundance ( $N$ ), contacts ( $C$ ), harvest ( $H$ ), release mortality ( $S$ ), dropoff mortality ( $D$ ), and impacts ( $I$ ) dependent on the contact rate ( $c$ ), the proportion of fish that are greater than or equal to the minimum size limit ( $p$ ), the release mortality rate ( $s$ ), and the dropoff mortality rate ( $d$ ). Because the model confines itself to age-3, for simplicity we have suppressed the use of a subscript denoting age for these quantities.

To project the cohort abundance forward in one month increments, total monthly impacts

$$I_t = \sum_{z,x} I_{tzx} \quad (6)$$

are first deducted from  $N_t$ , followed by application of the monthly natural mortality rate  $v$

$$N_{t+1} = (N_t - I_t) \times (1 - v). \quad (7)$$

Following projection of the cohort abundance across months, the age-3 impact rate ( $i_3$ ) is forecast as

$$i_3 = \frac{\sum_t I_t}{N_{\text{March}}}. \quad (8)$$

In practice,  $N_{\text{March}}$  is specified as 1.0, and  $i_3$  reduces to the numerator in (8).

In the following section, the submodels and input variables used to parameterize the  $c$ ,  $p$ ,  $s$ , and  $d$  rates are described.

## 4 Submodels and input variables

### 4.1 Contact rate

Age-3 month/area/sector contact rates have been estimated for years 2000–2009 through cohort reconstruction (O’Farrell et al. 2012). Pairing postseason estimates of  $c_{tzx}$  with postseason fishing effort estimates  $f_{tzx}$  allows for forecasting the contact rate per unit effort ( $\beta_{tzx}$ ), and ultimately, the expected contact rate in proposed fisheries.

Forecasts of  $\beta_{tzx}$  are determined by the slope of a zero-intercept linear model fitted to historical  $c_{tzx}$  and  $f_{tzx}$  data. Figures 3 and 4 displays these relationships for the commercial and recreational sectors, respectively. A ratio estimator is used to determine the month/area/sector contact rate per unit effort forecast,


$$\beta_{tzx} = \frac{\bar{c}_{tzx}}{\bar{f}_{tzx}} \quad (9)$$

following the methodology used for the KOHM (Mohr 2006a), where  $\bar{c}_{tzx}$  and  $\bar{f}_{tzx}$  denote the respective average of these quantities over the historical data.


Expected contact rates are then forecast using  $\beta_{tzx}$  and the effort forecast for that month/area/sector:

$$c_{tzx} = \beta_{tzx} \times f_{tzx}, \quad (10)$$

with the effort forecast determined as described in the following section.



**Figure 3.** Commercial sector age-3 contact rates, plotted as a function of fishing effort, by month and management area. Empty cells indicate strata with no data.



**Figure 4.** Recreational sector age-3 contact rates, plotted as a function of fishing effort, by month and management area. Empty cells indicate strata with no data.

## 4.2 Fishing effort

Fishing effort is not stock specific, and forecasts of fishing effort for each month/area/sector is made external to the WRHM. Effort forecasts are necessary inputs for the Klamath Ocean Harvest Model (KOHM) and Sacramento Harvest Model (SHM), as well as the WRHM, hence they are shared across models. Fishing effort forecast methods for both days-open fisheries and quota fisheries are documented in Mohr (2006a,b).

As described in Mohr (2006a), quota fishery effort is forecast in a different manner than days-open fishery effort. Effort expected in a quota fishery is determined by the size of the mixed-stock quota and the stock contribution rate of abundant target stocks (e.g., Klamath and Sacramento River fall Chinook). For quota fisheries occurring between September ( $y$ ) and February ( $y + 1$ ), the stock contribution rate of these target stocks is not known at the time of the PFMC preseason salmon management process because the ocean abundance of these fall run stocks has not yet been forecast (O’Farrell 2009). As a result, it is currently not possible to forecast quota fishery effort during this period using the customary method. If quota fisheries were planned in the SF and MO area during this period, fishing effort would need to be forecast using an alternative method. However, quota fisheries have very rarely occurred historically in the SF and MO area, and there is little reason to believe that they will be proposed in the near future.

## 4.3 Proportion legal size

Determination of  $p_{tzx}$  requires a specified minimum size limit ( $l_{tzx}^*$ ) and the mean ( $\mu_t$ ) and standard deviation ( $\sigma_t$ ) of the length distribution of age-3 SRWC in the ocean for month  $t$ . Minimum size limits are specified for nearly all ocean fisheries and are a standard input to the WRHM. The model used to estimate monthly size-at-age is described in O’Farrell et al. (2012, Appendix A). Size-at-age in month  $t$  is assumed to be normally distributed, so that given  $l_{tzx}^*$ ,  $\mu_t$ , and  $\sigma_t$ ,

$$p_{tzx} = P\{l \geq l_{tzx}^* | \mu_t, \sigma_t\} = 1 - \Phi(l_{tzx}^* | \mu_t, \sigma_t), \quad (11)$$

where  $P\{A\}$  denotes the probability of event  $A$ , and  $\Phi(\cdot)$  is the cumulative distribution function for the normal distribution.

#### 4.4 Release mortality rate

Based on the Salmon Technical Team (STT) review of hook and release mortality rates (STT 2000), we employ the conventional rate values of  $s_{tz(\text{com})} = 0.26$  for the commercial sector, and  $s_{tz(\text{rec})} = 0.14$  for the recreational sector when the method of fishing is exclusively trolling.

However, for the recreational sector, if the method of fishing known as “mooching” is used in addition to trolling in a particular month/area, then  $s_{tz(\text{rec})}$  is formulated as a weighted average of the troll release mortality rate (0.14) and the elevated mooch release mortality rate (0.422) (Grover et al. 2002). Mooching is a fishing technique that consists of drifting bait, which encourages the swallowing of the hook(s), and results in a higher proportion of gut-hooked fish relative to trolling (hence the high release mortality rate). Mooching has been popular in the SF and MO areas at times, but its use varies by month/area. Denoting  $\bar{\rho}_{tz}$  as the 5-year average of the month- and area-specific proportion of the recreational catch taken by mooching, the  $s_{tz(\text{rec})}$  forecast for the SF and MO areas is derived as

$$s_{tz(\text{rec})} = (\bar{\rho}_{tz} \times 0.422) + ((1 - \bar{\rho}_{tz}) \times 0.14). \quad (12)$$

Grover et al. (2002) presents details pertaining to the parameterization of this relationship.

#### 4.5 Dropoff mortality rate

Fish that contact fishing gear but are not brought to the boat may experience dropoff mortality. This source of mortality could result from a variety of causes, such as predation events or wounds inflicted by the fishing gear. Following STT (2000), we employ the conventional rate value of  $d = 0.05$ .



## 4.6 Natural mortality rate

The natural mortality annual rate is assumed to be 20 percent, and this corresponds to a monthly rate value of  $\nu = 0.0184$ . This is consistent with values used in the assessment of other Pacific salmon (e.g., Goldwasser et al. 2001; Mohr 2006a).

## 5 Discussion

We have formulated a harvest model capable of forecasting the annual age-3 impact rate for SRWC, given a proposed set of ocean salmon fishery management measures. This model is used as a tool to comply with the second component of the SRWC consultation standard, the age-3 impact rate control rule. The PFMC continues to have the customary fishery management controls of time/area/sector closures, spring/summer quotas, and minimum size limits available to meet the SRWC objectives.

Key inputs to the WRHM such as fishing effort and contact rates are based on relationships that utilize new information as it becomes available. Cohort reconstructions will be performed annually, providing new data that will be incorporated into the WRHM each year. This process allows the model to integrate changes in effort or exploitation patterns should they occur.

Contact rate forecasts are an important component of the WRHM. For the commercial sector, examination of contact rate and effort relationships illustrates the relative rarity of age-3, coded-wire tagged SRWC harvest. Fishery contacts are estimated by expanding the estimated harvest of coded-wire tagged SRWC by  $1/p_{t,zx}$ , with  $p_{t,zx}$  representing the proportion of SRWC expected to be of length greater than or equal to the minimal legal size (O'Farrell et al. 2012). In spring and early summer, a large proportion of age-3 SRWC are smaller than common commercial minimum size limits (O'Farrell et al. 2012, table A-2), hence landed catch is typically low. When landed catch is zero in a stratum, the contact rate is estimated to be zero as well, though it is possible that sublegal fish were contacted and released. For the commercial fishery, this interaction between SRWC size-at-age and the relatively large minimum size limits customary in this fishery likely contributes to the many instances of zero contact rates and the potential for contact rates to be underestimated in

cohort reconstructions. For the recreational fishery, there are many fewer instances of zero contact rates because age-3 SRWC become largely vulnerable to retention in the spring, and nearly all are vulnerable in the summer, given typical recreational sector size limits. As a result, fewer zero contact rate estimates exist, the potential for underestimation of contact rates in cohort reconstructions is low, and patterns in contact rates per unit effort are more clearly evident.

Since 2004, recreational fisheries south of Point Arena have been required to open no earlier than the first Saturday in April, while these fisheries traditionally opened in mid-February. Sufficient data do not exist to allow for robust contact rate estimation in February and March. Resumption of these early fisheries would result in highly uncertain forecasts of age-3 impact rates for those months because contact rates per unit effort ( $\beta_{t,x}$ ) would need to be assumed rather than directly estimated. Because of SRWC river return timing, it is likely that  $\beta_{t,x}$  is high in February and March (see Figure 11 in O'Farrell et al. 2012). This potential problem also exists for the commercial sector if fisheries prior to May 1 are proposed, and for both sectors if fisheries are proposed for late-fall or winter, when contact rate estimates are sparse or nonexistent. Because the first component of the SRWC consultation standard specifies retaining the time/area/fishery closure and size limit restrictions defined in 2004 (NMFS 2004), the expansion of salmon fisheries in the south of Point Arena area during these months is not currently being contemplated.

The WRHM shares many structural similarities to existing PFMC harvest models for Chinook salmon, and to the KOHM in particular. Like the KOHM, the WRHM is an age-structured model, though it only accounts for one age class. It is linked to a cohort reconstruction model with the same structure, which is updated annually. A size-at-age model is incorporated into both the KOHM and WRHM to allow for forecasting of release mortality incurred by sublegal size fish. Contact rates per unit effort are forecast in the same manner. Finally, many of the same conventions for  $s$ ,  $d$ , and  $v$  are shared across models. In contrast to the existing harvest models, the WRHM does not account for river fisheries, as SRWC are rarely harvested in the Sacramento River. Most importantly, neither preseason ocean abundance forecasts nor spawner escapement forecasts are made by the WRHM. As such, the WRHM can be considered a simplified harvest model in the same family as the KOHM and SHM.

## **6 Acknowledgments**

We wish to thank the PFMC Scientific and Statistical Committee Salmon Subpanel for their review of an earlier version of this document. In addition, we appreciate the reviews provided by William Satterthwaite and Valerie Brown of the Fisheries Ecology Division in Santa Cruz.

## References

- Fisher, F. W. (1994). Past and present status of Central Valley Chinook salmon. *Conservation Biology* 8, 870–873.
- Goldwasser, L., M. S. Mohr, A. M. Grover, and M. L. Palmer-Zwahlen (2001). The supporting databases and biological analyses for the revision of the Klamath Ocean Harvest Model. Unpublished report. National Marine Fisheries Service, Santa Cruz, CA.
- Grover, A. M., M. S. Mohr, and M. L. Palmer-Zwahlen (2002). Hook-and-release mortality of Chinook salmon from drift mooching with circle hooks: management implications for California's ocean sport fishery. In J. A. Lucy and A. L. Studholme (Eds.), *Catch and release in marine recreational fisheries*, pp. 39–53. American Fisheries Society, Bethesda, Maryland.
- Mohr, M. S. (2006a). The Klamath Ocean Harvest Model (KOHM): model specification. Unpublished report. National Marine Fisheries Service, Santa Cruz, CA.
- Mohr, M. S. (2006b). The Klamath Ocean Harvest Model (KOHM): parameter estimation. Unpublished report. National Marine Fisheries Service, Santa Cruz, CA.
- NMFS (2004). Supplemental Biological Opinion, on authorization of ocean salmon fisheries developed in accordance with the Pacific Coast Salmon Plan and proposed protective measures during the 2004 through 2009 fishing seasons as it affects Sacramento River Winter Chinook salmon. National Marine Fisheries Service, Southwest Region, Protected Resources Division.
- NMFS (2010). Authorization of ocean salmon fisheries pursuant to the Pacific Coast Salmon Fishery Management Plan and additional protective measures as it affects Sacramento River Winter Chinook Salmon. National Marine Fisheries Service, Southwest Region, Protected Resources Division.
- O'Farrell, M. R. (2009). Assessment of fall ocean Chinook salmon fisheries south of Cape Falcon, OR. Unpublished report. National Marine Fisheries Service, Santa Cruz, CA.
- O'Farrell, M. R., M. S. Mohr, A. M. Grover, and W. H. Satterthwaite (2012). Sacramento River winter Chinook cohort reconstruction: analysis of ocean fishery impacts. Report in preparation.

National Marine Fisheries Service, Santa Cruz, CA.

STT (2000). STT recommendations for hooking mortality rates in 2000 recreational ocean Chinook and coho fisheries. Pacific Fishery Management Council, 7700 NE Ambassador Place, Suite 101, Portland, Oregon 97220-1384.

## RECENT TECHNICAL MEMORANDUMS

SWFSC Technical Memorandums are accessible online at the SWFSC web site (<http://swfsc.noaa.gov>). Copies are also available from the National Technical Information Service, 5285 Port Royal Road, Springfield, VA 22161 (<http://www.ntis.gov>). Recent issues of NOAA Technical Memorandums from the NMFS Southwest Fisheries Science Center are listed below:

- NOAA-TM-NMFS-SWFSC-479 Passive acoustic beaked whale monitoring survey of the Channel Islands, CA. T.M. YACK, J. BARLOW, J. CALAMBOKIDIS, L. BALLANCE, R. PITMAN, and M. McKENNA  
(May 2011)
- 480 Determining transmitter drag and best-practice attachment procedures for sea turtle biotelemetry studies. T.T. JONES, B. BOSTROM, M. CAREY, B. IMLACH, J. MIKKELSEN, P. OSTAFICHUK, S. ECKERT, P. OPAY, Y. SWIMMER, J.A. SEMINOFF, and D.R. JONES  
(November 2011)
- 481 Ichthyoplankton, paralarval cephalopod, and station data for surface (Manta) and oblique (Bongo) plankton tows for California Cooperative Oceanic Fisheries Investigations Survey and California Current Ecosystem Survey cruises in 2008. W. WATSON and S.M. MANION  
(May 2011)
- 482 Toward a national animal telemetry observing network (ATN) for our oceans, coasts and great lakes: Workshop synthesis report. H. MOUSTAHDID, C GRIMES, J. KOCIK, B. BLOCK, K. HOLLAND, J. PAYNE, D. FOX, A. SEITZ, and C. ALEXANDER  
(July 2011)
- 483 Photographic catalog of California marine fish otoliths: Prey of California sea lions (*Zalophus californianus*). M.S. LOWRY  
(November 2011)
- 484 Effective strip widths for ship-based line-transect surveys of cetaceans. J. BARLOW, L.T. BALLANCE, and K.A. FORNEY  
(November 2011)
- 485 Fin Whale acoustics as a tool to assess stock structure in the North Pacific. B. JONES, S. RANKIN, and E. ARCHER  
(November 2011)
- 486 Spawning biomass of Pacific Sardine (*Sardinops sagax*) off U.S. in 2011. N.C.H. LO, B.J. MACEWICZ, and D.A. GRIFFITH  
(November 2011)
- 487 Assessment of the Pacific sardine resource in 2011 for U.S. management in 2012. K.T. HILL, P.R. CRONE, N.C.H. LO, B.J. MACEWICZ, E. DORVAL J.D. McDANIEL, and Y. GU  
(January 2012)
- 488 U.S. Pacific marine mammal stock assessments: 2011 J.V. CARRETTA, K.A. FORNEY, E. OLESON, K. MARTIEN, M.M. MOTO, M.S. LOWRY, J. BARLOW, J. BAKER, B. HANSON, D. LYNCH, L. CARSWELL, R.L. BROWNELL, JR., J. ROBBINS, D.K. MATTILA, K. RALLS, and M.C. HILL  
(April 2012)